



Constructing a Healthier Industry

Information Partner



what causes depression?

Most people assume that depression is caused simply by recent personal difficulties. Depression however, is often caused by the mix of recent events and other longer-term or personal risk factors.

Research indicates that ongoing difficulties, such as long-term unemployment or living in an abusive or uncaring relationship, are more likely to cause depression than recent life stressors. Depression can also run in families and some people will be at increased genetic risk. However, this doesn't mean that you will automatically become depressed if a parent or close relative has had the illness. Life circumstances are still likely to have an important influence on your chances of becoming ill.

It's also common for people to experience depression and anxiety at the same time.



Research indicates that ongoing difficulties, such as long-term unemployment or living in an abusive or uncaring relationship, are more likely to cause depression than recent life stressors.

It's important to note that you can't always identify the cause of depression nor change troubling circumstances. The most important thing is to recognise the depression and to seek help.

Remember, the sooner you get treatment, the greater the chance of a faster recovery.

The OzHelp Foundation

1300 OZHELP (1300 694 357)
www.ozhelp.org.au

BeyondBlue

For information about depression, anxiety and related drug and alcohol problems visit www.beyondblue.org.au or call the *beyondblue: the national depression initiative* info line on 1300 22 4636



Australian Government
Department of Health and Ageing

Common medical causes of depression include:


- Low thyroid function
- Brain injuries and diseases (eg. stroke, heart disease, head injury, epilepsy, Parkinson's Disease)
- Some forms of cancer
- Infectious diseases
- Blood vessel disease in the brain due to diabetes and/or hypertension
- Some steroid and hormonal treatments
- Chronic pain
- Quitting smoking.

High-risk personality being:

- A lifelong worrier
- A perfectionist
- Sensitive to personal criticism
- Unassertive
- Self-critical and negative
- Shy, socially anxious and having low self-esteem.

Common tests done by a doctor include:

- Full blood count and biochemistry
- Thyroid function tests
- Urine test for sugar and protein
- Occasionally, a brain scan.



Source: Hickie et al. Educational Health Solutions; 2000. Reproduced with permission www.spheregp.com.au.